



Integrated Management of Downy Brome in Winter Wheat

A PACIFIC NORTHWEST EXTENSION PUBLICATION • PNW668

Introduction

Downy brome (*Bromus tectorum* L.), also known as cheat-grass, was introduced into North America from the Mediterranean area of Europe. It was first identified in the eastern United States in 1861, and by 1914 this invasive weed had spread throughout the continent. Downy brome is adapted to climates with annual precipitation ranging from 6 to 22 inches. It can colonize both disturbed and undisturbed sites with a wide range of soil conditions.

Downy brome is a major weed problem in winter wheat (Figure 1). In eastern Washington, 54 downy brome plants per square foot reduced winter wheat yield by 92 percent (Rydrych and Muzik 1968). Downy brome is especially troublesome in low precipitation production areas where crop rotations are mostly limited to winter wheat followed by a year of summer fallow.

Downy brome is best controlled in wheat using integrated weed management (IWM). This approach involves a combination of management tools to reduce a weed population to an acceptable level while preserving the quality of natural resources.



Figure 1. Downy brome infestation in a winter wheat–fallow crop rotation. (Photo by Drew Lyon)

Identification

Downy brome is a winter annual grass that ranges from 12 to 24 inches in height at maturity. The seed head is 2–6 inches long, with 5–8 spikelet flowers (Figure 2). Magnification reveals greater detail of the ½ inch long seeds and their ½–¾ inch long awns (Figure 3), as well as the fine hairs on the leaf sheaths and blades (Figure 4).

Biology

Downy brome seed germination typically occurs in autumn shortly after the onset of rains when the soil temperature is about 70°F. Seeds can continue to germinate at soil temperatures between 35° and 40°F if soil moisture is adequate. Established plants overwinter in the vegetative stage, resume rapid growth in the early spring, and mature in May or June (4–6 weeks before winter wheat). Downy brome can also be produced from seeds germinating in the spring, although seed production is much more prolific from autumn-germinating plants. Newly mature downy brome seed requires a short after-ripening period for optimum germination. After-ripening, a process that



Figure 2. Mature downy brome plant showing slender, erect culms (tillers) supporting dense, slender, drooping panicles (seed heads) with spikelets of flowers. (Photo by Larry Burrill, formerly with Oregon State University)



Figure 3. Downy brome seeds with sharp-tipped, narrow awns. (Photo courtesy of Steve Hurst, USDA NRCS, Bugwood.org)



Figure 4. A downy brome leaf blade with sheath and ligule. The outgrowth from the top of the leaf sheath is membranous with frayed margins; it has no auricles (small ear-like projection from the base of many grass leaves). (Photo by Don Morishita)

many grasses must undergo after maturity before they can germinate, prevents seed from germinating prematurely under moist soil conditions.

Beyond this initial after-ripening period, little seed dormancy remains and germination occurs when favorable temperature and soil moisture conditions exist. Germination often exceeds 95 percent for seeds buried within 2 inches of the soil surface. Seeds buried below 4 inches may germinate but will rarely emerge above the soil surface. Seeds in aboveground crop or weed residue will survive longer than those seeds in direct contact with the soil. The majority of downy brome seeds remain viable in the soil for 2–3 years.

Downy brome exhibits rapid growth and development, making it highly competitive with winter wheat, particularly when the timing of emergence for the two is similar. The primary root system, which develops from the seed, grows throughout the fall and winter at soil temperatures just above freezing. Secondary or adventitious roots emerging from the plant crown are usually initiated in the fall or winter and become well developed before winter wheat

resumes growth in the spring. The finely divided fibrous root system is highly efficient in exploiting soil moisture and nutrients. While some roots can reach 4 feet deep, more than 90 percent of the root mass is contained in the top 15 inches of soil.

By the time winter wheat begins rapid spring growth, downy brome plants are often tillering or jointing. Heading usually occurs 4 weeks earlier than winter wheat. Seeds mature by late spring and seed shatter often begins before wheat harvest. Under ideal environmental conditions, (represented by a dense infestation), downy brome can produce over 500 pounds of seed per acre, resulting in the potential for 125 million plants.

Management

Using IWM for downy brome in winter wheat is centered on preventing the weed from becoming established in new areas, growing a competitive crop, and keeping weeds off-balance by changing management practices.

All the practices discussed here can be effective against downy brome, provided the associated cautions are taken. Field burning, commonly used in the past, is not considered part of IWM for downy brome because the many undesirable consequences far outweigh the questionable benefits.

Elimination of seed sources

Because downy brome is a prolific seed producer, eliminating seed sources can be an effective preventive control strategy. However, this often requires sustained effort over time. Apply a combination of the following methods that fit best:

- Plant crops using clean, certified seed.
- Seed perennial, cool-season grasses in waste areas and field borders. Vigorous stands of grasses or grass–legume combinations are highly competitive with downy brome and other annual weeds. (See the sidebar on grass options for suggestions.)

Perennial, cool-season grass options for waste areas and field borders to prevent downy brome infestation.

Precipitation zone (inches/year)	Grass cultivar
High (20+)	'Durar' hard fescue 'Rush' intermediate wheatgrass 'Alkar' tall wheatgrass
Intermediate (15–20)	'Durar' hard fescue 'Nordan' crested wheatgrass 'Bannock' thickspike wheatgrass
Low (less than 15)	'Covar' sheep fescue 'Sodar' streambank wheatgrass 'Vavilov II' Siberian wheatgrass

- Destroy downy brome seedlings in cultivated fields before they produce seed.
- Control small patches or area infestations before they spread.
- Where perennial grass borders are not feasible, consider a) using herbicides or tillage to kill downy brome and b) cropping field borders and roadside ditches.
- Use herbicides that do not kill established perennial grasses around field borders.
- Mow small infestations close to the ground in pastures, roadsides, and waste areas where cultivation or herbicides are not feasible. Plan to complete mowing before viable downy brome seeds are produced. More than one mowing may be necessary to prevent tillers from producing seed.

Crop diversity

Crop rotation is one of the most effective control measures for downy brome because of the relatively short life span (2–3 years) of the seed in soil. The highest rate of success is associated with lengthening the period between winter wheat crops by including 2 or more years of various combinations of spring-planted crops with or without fallow.

Broadleaf crops, such as dry pea, chickpea, lentils, and canola, are good choices for rotation with winter wheat because they allow use of Group 1 herbicides (ACC-ase inhibitors), such as sethoxydim (Poast), clethodim (Select Max), or quizalofop (Assure II, Targa), to manage seedling downy brome. These same herbicides can be used in many irrigated broadleaf crops grown in rotation with winter wheat. In lower precipitation areas, possible crop rotations are 1) winter wheat–summer fallow–winter canola or spring wheat–summer fallow, 2) winter wheat–spring barley or spring canola–summer fallow, and 3) winter wheat–spring barley or spring canola–spring wheat–summer fallow.

In higher precipitation areas with annual cropping, rotations including 2 years of spring crops such as barley, canola, chickpea, lentils, or dry pea between winter wheat crops can effectively reduce downy brome populations. These types of rotations allow for fall and/or spring control of germinating downy brome over 2 or more years. This can prevent new seed production and deplete the downy brome seed reserve in the soil.

The later in the spring these crops can be planted, the more effective they will be for controlling downy brome. Later planting allows more time for downy brome emergence in the spring and a higher percentage of weed seedlings to be killed with nonselective herbicides or tillage before planting. In addition, late-emerging downy brome may not be exposed to enough cold to vernalize (that is, become reproductive).

A significant disadvantage of delayed spring planting is potentially lower crop yield. Delayed planting tends to have a greater negative effect on yields for spring wheat yields than spring barley, which may make barley a better

choice for use in crop rotations to control downy brome. However, there are fewer herbicide options for controlling downy brome in barley than in spring wheat. Chickpeas may be the best alternative because it is generally planted later than both of these spring small grains and allows selective Group 1 grass herbicides to be used.

The key aspect of crop rotation for control of downy brome is to prevent any new seed production for at least 2 years between winter wheat crops. During this time, the seed bank will decline significantly as the result of germination, predation, and other forms of seed mortality.

Fallow management

Germination and subsequent control of brome seedlings are critical to good fallow management. This approach begins with even distribution of the wheat crop residues during crop harvest. Downy brome seed will be spread with the chaff and finer residue particles, allowing for seed-to-soil contact to encourage germination. Straw choppers on combines help spread the straw and make it easier to get good seed-to-soil contact later. Since rotary harvesters break up the straw, choppers are not usually needed.

After the crop residues have been adequately distributed, consider non-inversion tillage to “plant” downy brome seed for optimum germination during the fallow period. This is most useful in winter wheat–fallow rotations where there is only one year between winter wheat crops to deplete the soil seed bank. Research conducted in eastern Washington in a winter wheat–fallow rotation found that using a sweep plow or disc after wheat harvest resulted in fewer downy brome plants in the following winter wheat crop than using a skew treader or no tillage after harvest (Young et al. 2014) (Table 1).

Use non-inversion tillage or a short-residual herbicide such as glyphosate in the fall following downy brome emergence to control seedlings. Herbicides are more effective than tillage when soil is moist and plants are actively growing. Unlike tillage, herbicides do not destroy additional crop residues. For tillage to work well, the soil must be dry and the air temperature must be warm enough (above 80°F) to desiccate plants within about 30 minutes.

During the spring of the fallow year, it is important not to allow any new downy brome seed production to occur.

Table 1. Downy brome densities in winter wheat following 5 postharvest tillage treatments (amended from Young et al. 2014).

Treatment	Depth of tillage (inches)	Average number of plants/ft ²
No-till	—	17a*
Skew treader	1-2	14a
Harrow	1	13ab
Sweep plow	4-5	10b
Disk	3-4	10b

*Treatment averages followed by the same letter are not significantly different from each other at the 95% confidence level.

This can be difficult, particularly with prolonged wet conditions, because of the rapid development of downy brome from heading through seed set. Pollination occurs very quickly after heading and is difficult to recognize. Once pollination has occurred, at least some viable seed will be produced even if the plant is subsequently killed by tillage or herbicides. Research conducted in Oregon and Washington found that downy brome typically produces viable seed when approximately 1800 growing degree days (GDD; 32°F base temperature) have accumulated from January 1 (Ball et al. 2004). Table 2 provides some average dates for attaining 1800 GDD at locations across the inland Pacific Northwest.

Plowing

If soil erosion and conservation compliance are not problems, a moldboard plow may be used to bury downy brome seeds at least 4 inches deep. This can provide 95 percent control of downy brome. Subsequent plowing should be avoided for at least 4 years to prevent viable seed from returning to the soil surface. Plowing should be restricted to areas with the greatest downy brome infestations to limit negative effects on soil health, including erosion.

Fertilizer

Downy brome responds dramatically to nitrogen fertilization of winter wheat. Surface-applied nitrogen, either as commercial fertilizer or manure, has been shown to triple downy brome height and seed yield. Test soil and apply only the recommended amount of nitrogen fertilizer. Over-application can result in crop yield reduction due to downy brome competition.

Table 2. Average dates when 1800 growing degree days have accumulated at several locations in Washington, Oregon, and Idaho.

Location	Date*
Washington	
Walla Walla	May 16
Lind	June 6
Pullman	June 18
Davenport	June 25
Oregon	
Corvallis	May 11
Pendleton	May 18
Moro	June 8
La Grande	June 13
Idaho	
Lewiston	May 19
Twin Falls	June 13
Pocatello	June 18
Idaho Falls	June 22

*Downy brome plants must be killed prior to these dates to ensure no new seed is produced.

Deep banding of nitrogen fertilizer improves winter wheat yield and crop competitiveness over downy brome compared to a broadcast application. Deep band nitrogen fertilizer early in the fallow season rather than at or just prior to planting. Do not spring topdress nitrogen fertilizer in winter wheat fields with downy brome infestations, as the crop rarely benefits and the practice increases weed seed production and weed water use.

If phosphorus fertilizer is needed, a deep band or starter placement can stimulate wheat root growth and increase crop competitiveness with downy brome. However, there is often less crop response to phosphorus fertilizer in early seeded fields.

Timing of winter wheat seeding

Seeding winter wheat at the optimum date for the area is important to control downy brome. Most downy brome will emerge from seed within the top ½ inch of the surface but won't germinate until moisture is sufficient. Seeding through a dry mulch layer into a moist seed zone with a deep furrow drill will allow wheat to emerge before downy brome and improve its competitive advantage. Research has shown that wheat emerging 3 weeks prior to downy brome will prevent significant yield losses from light to moderate infestations of about 10 plants per square foot or less (Stahlman and Miller 1990). Yield loss increases dramatically when downy brome emerges within 7–10 days of wheat. Early winter wheat emergence also allows the differential growth between wheat and downy brome that improves the efficacy and crop safety of postemergence herbicides.

If sufficient rain (generally considered to be more than a tenth of an inch) occurs just before the anticipated winter wheat planting, the best strategy to control downy brome is to delay seeding until the weed has emerged and either been tilled or sprayed with a nonselective herbicide. However, delaying seeding beyond the optimum seeding date may reduce fall crop growth, competitiveness with downy brome, and yield. If rain shortly after planting results in soil crusting and the need to replant, wait until downy brome has emerged so that it can be killed with herbicides or tillage prior to replanting winter wheat. With later seeding in moist conditions, winter wheat and downy brome will emerge at the same time, resulting in greater weed competition and reduced efficacy or increased crop injury from postemergence herbicides.

If downy brome infestation is severe, consider substituting a spring crop in place of winter wheat in the rotation. (See Crop diversity discussion above.)

Herbicides

Several herbicides are labeled for selective control or suppression of downy brome in winter wheat. Some of the most effective of these are Group 2 (ALS-inhibitors) herbicides such as sulfosulfuron (Maverick, Outrider), propoxycarbazone (Olympus), and pyroxsulam (PowerFlex HL). These products provide excellent control of downy brome when applied in the fall, but are more inconsistent with

spring applications. They pose little risk for injuring winter wheat.

Beyond is another Group 2 herbicide that provides excellent control of downy brome when properly applied, but it is specific to Clearfield wheat varieties that contain the gene or genes that confer tolerance to imazamox. Tolerance means that the winter wheat variety with the gene(s) is able to withstand a recommended rate of Beyond with minimal risk of crop injury. Wheat varieties that do not contain this gene are either killed or seriously injured by Beyond.

There is some concern that downy brome will develop resistance to these Group 2 herbicides, so they should not be used repeatedly. Herbicides with other mechanisms of action should be rotated with Group 2 herbicides. Axiom provides very good to excellent control of downy brome that has not yet emerged. It contains two active ingredients, flufenacet (Group 15) and metribuzin (Group 5). Zidua contains pyroxasulfone (Group 15) and provides control of downy brome similar to Axiom. Crop injury is a potential concern with Axiom and Zidua, so it is critical to consult these labels for planting and application restrictions.

Other herbicides labeled for use in winter wheat, such as Far-Go (Group 8), metribuzin, metribuzin + Finesse (Groups 5 and 2), and Treflan (Group 3), can be used to suppress downy brome but generally do not provide the same level of control as the previously mentioned products.

For current herbicide control strategies for downy brome, refer to the *Pacific Northwest Weed Management Handbook* and contact local county Extension educators or agricultural professionals. As with all crop protection chemicals, read and follow label directions and understand their proper use.

Acknowledgements

This publication updates and replaces Nesse and Ball, *Downy Brome* (Oregon State University, PNW474, 1994); and Yenish et al., *Managing Downy Brome under Conservation Tillage Systems in the Inland Northwest Cropping Region* (Washington State University, PNW509, 1998). The authors of PNW668 recognize the contributions made by the authors of these two previous publications:

Author list for *Downy Brome* (PNW474):

- Philip Nesse, OSU Extension agent, Gilliam/Morrow Counties, OR
- Daniel Ball, OSU Weed Scientist, Columbia Basin Agricultural Research Center, Pendleton, OR

Author list for *Managing Downy Brome under Conservation Tillage Systems in the Inland Northwest Cropping Region* (PNW509):

- Joe Yenish, WSU Extension Weed Scientist, Pullman, WA

- Roger Veseth, WSU/UI Extension Conservation Tillage Specialist, Moscow, ID
- Alex Ogg, USDA-ARS Plant Physiologist, Nonirrigated Weed Science Research Unit, Pullman, WA
- Donn Thill, UI Weed Scientist, Moscow, ID
- Dan Ball, OSU Weed Scientist, Pendleton, OR
- Frank Young, USDA-ARS Research Agronomist, Nonirrigated Weed Science Research Unit, Pullman, WA
- Eric Gallandt, WSU Weed Scientist, Pullman, WA
- Don Morishita, UI Extension Weed Scientist, Twin Falls, ID
- Carol Mallory-Smith, OSU Weed Scientist, Corvallis, OR
- Don Wysocki, OSU Extension Soil Scientist, Pendleton, OR
- Tom Gohlke, NRCS State Agronomist, Portland, OR

References

- Ball, D.A., S.M. Frost, and A.I. Gitelman. 2004. Predicting Timing of Downy Brome (*Bromus tectorum*) Seed Production Using Growing Degree Days. *Weed Science* 52: 518–524.
- Peachey, E., ed. (updated annually). *Pacific Northwest Management Handbook*. Oregon State University. <http://pnwhandbooks.org/weed/>.
- Rydrich, D.J. and T.J. Muzik. 1968. Downy Brome Competition and Control in Dryland Wheat. *Agronomy Journal* 60: 279–280.
- Stahlman, P.W., and S.D. Miller. 1990. Downy Brome (*Bromus tectorum*) Interference and Economic Thresholds in Winter Wheat (*Triticum aestivum*). *Weed Science* 38: 224–228.
- Young, F.L., A.G. Ogg, Jr., and J.R. Alldredge. 2014. Postharvest Tillage Reduces Downy Brome (*Bromus tectorum* L.) Infestations in Winter Wheat. *Weed Technology* 28: 418–425.

By **Drew J. Lyon**, Extension Small Grains Weed Scientist, Washington State University; **Andrew G. Hulting**, Extension Weed Science Specialist, Oregon State University; **Don W. Morishita**, Extension Weed Science Specialist, University of Idaho; and **Frank L. Young**, Research Agronomist, USDA-ARS, Pullman, WA.

Use pesticides with care. Apply them only to plants, animals, or sites as listed on the label. When mixing and applying pesticides, follow all label precautions to protect yourself and others around you. It is a violation of the law to disregard label directions. If pesticides are spilled on skin or clothing, remove clothing and wash skin thoroughly. Store pesticides in their original containers and keep them out of the reach of children, pets, and livestock.

Pacific Northwest Extension publications are produced cooperatively by the three Pacific Northwest land-grant universities: Washington State University, Oregon State University, and the University of Idaho. Similar crops, climate, and topography create a natural geographic unit that crosses state lines. Since 1949, the PNW program has published more than 650 titles, preventing duplication of effort, broadening the availability of faculty specialists, and substantially reducing costs for the participating states.

Pacific Northwest Extension publications contain material written and produced for public distribution. You may reprint written material, provided you do not use it to endorse a commercial product. Please reference by title and credit Pacific Northwest Extension publications.

Copyright 2015 Washington State University.

Order information:

Washington State University Extension
<http://pubs.wsu.edu>
Fax 509-335-3006
Toll-free phone 800-723-1763
ext.pubs@wsu.edu

Oregon State University Extension Service
<http://extension.oregonstate.edu/catalog>
Fax 541-737-0817
Toll-free phone 800-561-6719
[puborders@oregonstate.edu](mailto:pубorders@oregonstate.edu)

University of Idaho Extension
<http://www.cals.uidaho.edu/edComm/catalog.asp>
Fax 208-885-4648
Phone 208-885-7982
calspubs@uidaho.edu

Published and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914, by Washington State University Extension, Oregon State University Extension Service, University of Idaho Extension, and the U.S. Department of Agriculture cooperating. WSU Extension programs, activities, materials, and policies comply with federal and state laws and regulations on nondiscrimination regarding race, sex, religion, age, color, creed, and national or ethnic origin; physical, mental, or sensory disability; marital status or sexual orientation; and status as a Vietnam-era or disabled veteran. Washington State University Extension, Oregon State University Extension Service, and University of Idaho Extension are Equal Opportunity Employers. Evidence of noncompliance may be reported through your local Extension office. Trade names have been used to simplify information; no endorsement is intended. Published April 2015. *This publication replaces PNW474 and PNW509.*